


概述 (Summary)

英文全称	Carboxylesterase 5/CES5
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Carboxylesterase 2E is produced by our Mammalian expression system and the target gene encoding Gln27-His556 is expressed with a 6His tag at the C-terminus.
Accession #	Q8BK48
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	60.2 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse CES2E (C-6His)
Catalog #: PHM0225



背景 (Background)

分子别名 (Alternative Names)

Pyrethroid hydrolase Ces2e; carboxylesterase 2E; Ces5

背景介绍 (References)

Carboxylesterase 5 (CES5), also called cauxin or CES7, is a member of carboxylesterases family which plays an important role in the hydrolysis of ester and amide bonds. Carboxylesterase is a type of enzyme that capable of hydrolyzing a variety of carboxylic acid esters and its widely distributed in cells especially in mammalian liver. CES5 is with broad substrate specificity ranging from small molecule esters to longchain fatty acid esters and thioesters. It has been previously reported CES5 was in high concentrations in the urine (cauxin) of adult male cats, and within a protein complex of mammalian male epididymal fluids. Roles for CES5 may include regulating urinary levels of male cat pheromones, catalyzing lipid transfer reactions within mammalian male reproductive fluids, and protecting neural tissue from drugs and xenobiotics.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.