


概述 (Summary)

英文全称	Ectodysplasin A2 Receptor/EDA2R/TNFRSF27
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Ectodysplasin A2 Receptor is produced by our Mammalian expression system and the target gene encoding Met1-Thr138 is expressed with a 6His tag at the C-terminus.
Accession #	Q8BX35
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	16.4 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse EDA2R (C-6His)
Catalog #: PHM0550



背景 (Background)

分子别名 (Alternative Names)

Ectodysplasin A2 receptor; EDA-A2 receptor; EDA-A2R; Tumor necrosis factor receptor superfamily member XEDAR; Tumor necrosis factor receptor superfamily member 27; X-linked ectodysplasin-A2 receptor; EDAA2R; TNFRSF27; XEDAR; EDAR2

背景介绍 (References)

Tumor necrosis factor receptor superfamily member 27, also known as XEDAR and EDA2R, is a type III transmembrane protein of the TNFR superfamily. EDA2R consists of extracellular domain (ECD) with 3 cysteine-rich repeats and a single transmembrane domain but lacks an N-terminal signal peptide. EDA2R is widely expressed, notably in embryonic basal epidermal cells and maturing hair follicles. Even though it does not contain a cytoplasmic death domain, EDA2R can associate with Fas and induce EDA-A2 dependent apoptosis. Its transcription is directly induced by p53, and it mediated cell death is p53 dependent. It is down-regulated in breast, colon, and lung cancers, particularly in cases with p53 mutations. It also plays a role in EDA-A2 induced skeletal muscle degeneration and osteoblast differentiation. Mutations in the EDA gene are associated with the X-linked form of Hypohidrotic Ectodermal Dysplasia (HED), a disease typically characterized by abnormal hair, teeth and sweat glands.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.