


概述 (Summary)

英文全称	Exostosin-Like 2/EXTL2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Exostosin-like 2 is produced by our Mammalian expression system and the target gene encoding Asn43-Met330 is expressed with a 6His tag at the N-terminus.
Accession #	Q9ES89
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	33.6 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse EXTL2 (N-6His)
Catalog #: PHM0609



背景 (Background)

分子别名 (Alternative Names)

Exostosin-like 2; Extl2; Alpha-1;4-N-acetylhexosaminyltransferase EXTL2; Alpha-GalNAcT EXTL2; EXT-related protein 2; Glucuronyl-galactosyl-proteoglycan 4-alpha-N-acetylglucosaminyltransferase

背景介绍 (References)

Exostosin-like 2 (EXTL2) is a member of the exostosin (EXT)-related family which contains five members: EXT1, EXT2, EXTL1, EXTL2, and EXTL3. Studies have shown that EXT gene family members have the activities of heparan sulfate-synthesizing glycosyltransferases. EXT1 and EXT2, which have been identified as causal genes for hereditary multiple exostoses, have HS-GlcAT-II and GlcNAcT-II activities. EXTL1 has GlcNAcT-II activity and EXTL3 has GlcNAcT-I and -II activities. EXTL2 has GlcNAcT-I and N-acetylgalactosaminyltransferase activities, and transfers a GlcNAc residue to the tetrasaccharide linkage region when this region is phosphorylated by a xylose kinase 1 (FAM20B) and thereby terminate chain elongation. In mice, lack of EXTL2 causes glycosaminoglycan (GAG) overproduction and structural changes of GAGs associated with pathological processes.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.