


概述 (Summary)

英文全称	CD95/TNFRSF6/FAS
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Apoptosis-mediating Surface Antigen FAS is produced by our Mammalian expression system and the target gene encoding Gln22-Arg169 is expressed with a 6His tag at the C-terminus.
Accession #	P25446
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	17.4 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse Fas (C-6His)
Catalog #: PHM0624



背景 (Background)

分子别名 (Alternative Names)

Tumor necrosis factor receptor superfamily member 6; Apo-1 antigen; Apoptosis-mediating surface antigen FAS; FASLG receptor; CD95; Fas; TNFRSF6

背景介绍 (References)

Mouse Apoptosis-mediating surface antigen FAS (Fas) belongs to the death receptor subfamily of the TNF receptor superfamily and is designated TNFRSF6. Mouse Fas contains 1 death domain and 3 TNFR-Cys repeats. It detected in various tissues including thymus, liver, lung, heart, and adult ovary. As a receptor for TNFSF6/FASLG, The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases mediating apoptosis. FAS-mediated apoptosis may have a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.