


概述 (Summary)

英文全称	Fc gamma RIII/CD16/FCGR3
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Fc Gamma Receptor III is produced by our Mammalian expression system and the target gene encoding Leu32-Thr215 is expressed with a 6His tag at the C-terminus.
Accession #	Q5D5I8
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	21.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse CD16 (C-6His)
Catalog #: PHM0631



背景 (Background)

分子别名 (Alternative Names)

Low affinity immunoglobulin gamma Fc region receptor III; Fcgr3; Fc gamma Receptor III; CD-antigen 16; CD16; FcRIII; IgG Fc receptor III

背景介绍 (References)

Low affinity immunoglobulin gamma Fc region receptor III (Fc gamma RIII/CD16) is a member of the Ig superfamily. Based on close relationships in their extracellular domains, the Fc gamma Rs have been divided into three classes composing of Fc gamma RI (CD64), Fc gamma RII (CD32), and Fc gamma RIII (CD16). Each group may be encoded by multiple genes and exist in different isoforms depending on species and cell type. Mouse CD16 is a type I transmembrane protein having two extracellular Ig-like domains consisting of immunoglobulin domain, repeat, signa and transmembrane, transmembrane helix. It is expressed on a variety of myeloid and lymphoid cells and associates with Fc R gamma to deliver an activating signal upon ligand binding. Fcgr3 is IgG binding and activation or inhibition of immune responses such as antibody-dependent cellular cytotoxicity, phagocytosis, cell surface receptor signaling pathway and positive regulation of type I/IIa/III hypersensitivity.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.