

产品名称: Recombinant Mouse G-CSF (C-6His)
产品货号: PHM0758




概述 (Summary)

英文全称	G-CSF/CSF1/Granulocyte Colony-Stimulating Factor
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<0.01 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Granulocyte Colony-Stimulating Factor is produced by our Mammalian expression system and the target gene encoding Val31-Ala208 is expressed with a 6His tag at the C-terminus.
Accession #	P09920
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	19.8 KDa
制剂 (Form)	Lyophilized from a 0.2 μm filtered solution of 10mM Sodium Citrate, 0.1% Tween 20, pH 3.5.
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

产品名称: Recombinant Mouse G-CSF (C-6His)
产品货号: PHM0758



背景 (Background)

分子别名 (Alternative Names)

Granulocyte colony-stimulating factor; Csf3; G-CSF

背景介绍 (References)

Granulocyte colony-stimulating factor (G-CSF) is a growth factor and an essential cytokine which belongs to the IL-6 superfamily. Granulocyte/macrophage colony-stimulating factors are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages. G-CSF binding to its receptor G-CSF-R which belongs to the cytokine receptor type I family depends on the interaction of alpha-helical motifs of the former and two fibronectin type III as well as an immunoglobulin-like domain of the latter. G-CSF is a cytokine that have been demonstrated to improve cardiac function and perfusion in myocardial infarction. And it was initially evaluated as a stem cell mobilizer and erythropoietin as a cytoprotective agent.

注意事项 (Note)

For research use only .