


概述 (Summary)

英文全称	Growth Hormone Receptor/GHR/GHBP
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Growth Hormone Receptor is produced by our Mammalian expression system and the target gene encoding Met1-Gln273 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q3UP14
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	58.3 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse GHR (C-Fc)
Catalog #: PHM0763



背景 (Background)

分子别名 (Alternative Names)

growth hormone binding protein; growth hormone receptor; serum binding protein; Somatotropin receptor;GHR; GH receptor; GHBP

背景介绍 (References)

Growth hormone receptor is a transmembrane receptor for growth hormone (GH). GH is a single-chain polypeptide that is mainly synthesized and released from the anterior pituitary gland and plays essential roles in growth, development and metabolism. GH exerts its physiological actions via GH binding to its receptor in its extracellular domain. Binding of growth hormone to the receptor leads to receptor dimerization and the activation of an intra- and intercellular signal transduction pathway leading to growth. Growth hormone receptor has been shown to interact with SGTA, PTPN11, Janus kinase 2, Suppressor of cytokine signaling 1 and CISH.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.