


概述 (Summary)

英文全称	IGFBP-6/IGF Binding Protein 6
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Insulin-Like Growth Factor-Binding Protein 6 is produced by our Mammalian expression system and the target gene encoding Ala26-Gly238 is expressed with a 6His tag at the C-terminus.
Accession #	P47880
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	23.7 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)



背景 (Background)

分子别名 (Alternative Names)

Insulin-like growth factor-binding protein 6; IBP-6; IGF-binding protein 6; IGFBP-6; Igfbp6; IBP6; IGF binding protein 6; insulin-like growth factor-binding protein 6

背景介绍 (References)

Insulin-like growth factors (IGFs) comprise a family of endocrine, paracrine and autocrine polypeptides consisting of the ligands IGF1 and IGF2, two receptors (IGF1R, IGF2R), at least 6 IGF-binding proteins (IGFBPs) and IGFBP proteases. Among the binding proteins, IGFBP6 is unique because of its N-terminal disulfide linkages and its marked binding preference for IGF2. It is a potent inhibitor of the interaction between IGF2 and its receptor IGF1R, thus preventing major functions of IGF2, such as induction of proliferation, differentiation, cell adhesion, or colony formation. In particular, IGFBP-6 inhibited the growth of neuroblastoma and rhabdomyosarcoma xenografts. GFBP-6 is expressed in many tissues, including lung, liver, gut and the central nervous system.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.