


## 概述 (Summary)

<b>英文全称</b>	IL-13/Interleukin-13 (Pro22-Phe131)
<b>纯度 (Purity)</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>内毒素 (Endotoxin level)</b>	<1 EU/μg as determined by LAL test.
<b>蛋白构建 (Construction)</b>	Recombinant Mouse Interleukin-13 is produced by our Mammalian expression system and the target gene encoding Pro22-Phe131 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	P20109
<b>蛋白标签 (Tag)</b>	
<b>表达宿主 (Host)</b>	Human Cells
<b>种属 (Species)</b>	Mouse
<b>预测分子量 (Predicted MW)</b>	13.1 KDa
<b>蛋白形态 (Form)</b>	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
<b>储存缓冲液 (Buffer)</b>	
<b>运输方式 (Shipping)</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>稳定性&amp;储存 (Stability &amp;Storage)</b>	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
<b>复溶 (Reconstitution)</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## 电泳图 (SDS-PAGE image)



## 背景 (Background)

### 分子別名 (Alternative Names)

Interleukin-13; IL-13; T-Cell Activation Protein P600; IL13; IL-13

### 背景介绍 (References)

Mouse interleukin 13 (mIL-13) is a pleiotropic cytokine produced by activated Th2 cells. IL-13 induces B cell proliferation and immunoglobulin production. It contains a four helical bundle with two internal disulfide bonds. Mouse IL13 shares 58% sequence identity with human protein and exhibits cross-species activity. IL13 signals via receptor IL13R (type2, IL4R) and activates STAT-6. IL13 initially binds IL-13R $\alpha$ 1 with low affinity and triggers association of IL4R $\alpha$ , generating a high affinity heterodimeric receptor IL13R and eliciting downstream signals. IL13 also binds IL-13R $\alpha$ 2 with high affinity, which plays a role in a negative feedback system of IL13 signaling. IL13 is an important mediator of allergic inflammation and disease.

## 注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.