


概述 (Summary)

| | |
|-----------------------------|---|
| 英文全称 | IL-13/Interleukin-13 (Ser26-Phe131) |
| 纯度 (Purity) | Greater than 95% as determined by reducing SDS-PAGE |
| 内毒素 (Endotoxin level) | <1 EU/μg as determined by LAL test. |
| 蛋白构建 (Construction) | Recombinant Mouse Interleukin-13 is produced by our Mammalian expression system and the target gene encoding Ser26-Phe131 is expressed with a 6His tag at the C-terminus. |
| Accession # | P20109 |
| 蛋白标签 (Tag) | |
| 表达宿主 (Host) | Human Cells |
| 种属 (Species) | Mouse |
| 预测分子量 (Predicted MW) | 12.7 KDa |
| 蛋白形态 (Form) | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4. |
| 储存缓冲液 (Buffer) | |
| 运输方式 (Shipping) | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. |
| 稳定性&储存 (Stability &Storage) | Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months. |
| 复溶 (Reconstitution) | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

电泳图 (SDS-PAGE image)



背景 (Background)

分子別名 (Alternative Names)

Interleukin-13; IL-13; T-Cell Activation Protein P600; IL13; IL-13

背景介绍 (References)

Mouse interleukin 13 (mIL-13) is a pleiotropic cytokine produced by activated Th2 cells. IL-13 induces B cell proliferation and immunoglobulin production. It contains a four helical bundle with two internal disulfide bonds. Mouse IL13 shares 58% sequence identity with human protein and exhibits cross-species activity. IL13 signals via receptor IL13R (type2, IL4R) and activates STAT-6. IL13 initially binds IL-13R α 1 with low affinity and triggers association of IL4R α , generating a high affinity heterodimeric receptor IL13R and eliciting downstream signals. IL13 also binds IL-13R α 2 with high affinity, which plays a role in a negative feedback system of IL13 signaling. IL13 is an important mediator of allergic inflammation and disease.

注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.