


概述 (Summary)

英文全称	IL-2 R alpha/CD25/IL2-RA/IL-2R α/IL-2 receptor subunit alpha
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Interleukin-2 Receptor Subunit Alpha is produced by our Mammalian expression system and the target gene encoding Glu22-Lys236 is expressed with a 6His tag at the C-terminus.
Accession #	P01590
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	25.5 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse IL-2RA (C-6His)
Catalog #: PHM0914



背景 (Background)

分子别名 (Alternative Names)

CD25; IL-2 R alpha; IL2RA; CD25 antigen; IDDM10; IL-2 receptor subunit alpha; IL2R; IL-2R subunit alpha; IL-2-RA; IL2-RA; interleukin 2 receptor; alpha; interleukin-2 receptor subunit alpha; p55; TAC antigen; TCGFR

背景介绍 (References)

The interleukin 2 receptor alpha (IL2RA), also called CD25, is a type I transmembrane protein that presents on activated T cells, activated B cells, some thymocytes, myeloid precursors, and oligodendrocytes. IL2RA is a member of cytokine receptors family that utilizes the common gamma chain subunit (γ c). IL2RA is expressed in most B-cell neoplasms, some acute nonlymphocytic leukemias, neuroblastomas, and tumor infiltrating lymphocytes. IL2RA associates with IL2RB (CD122) to form a heterodimer that can act as a high-affinity receptor for IL-2.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.