产品名称：Recombinant Mouse IL-23
产品货号：PHM1001

概述（Summary）

英文全称	IL-23/Interleukin-23(IL23A&IL12B)
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	Recombinant Mouse Interleukin-23 is produced by our Mammalian expression system and the target gene encoding Val22-Ala196&Met23-Ser335 is expressed.
Accession #	Q9EQ14&P43432
表达宿主（Host）	Human Cells
种属（Species）	Mouse
预测分子量（Predicted MW）	19.7&35.8 KDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
运输方式（Shipping）	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图（SDS-PAGE image）

产品名称：Recombinant Mouse IL-23
产品货号：PHM1001



背景（Background）

分子别名（Alternative Names） SGRF; IL-23p19; CLMF p40; IL-12 subunit p40; NKSF2

背景介绍（References） Interleukin 23 (IL-23) is a heterodimeric cytokine composed of two disulfide-linked subunits, a p19 subunit that is unique to IL-23, and a p40 subunit that is shared with IL-12. The p19 subunit has homology to the p35 subunit of IL-12, as well as to other single chain cytokines such as IL-6 and IL-11. The p40 subunit is homologous to the extracellular domains of the hematopoietic cytokine receptors. Although p19 is expressed by activated macrophages, dendritic cells, T cells, and endothelial cells, only activated macrophages and dendritic cells express p40 concurrently to produce IL-23. IL-23 has biological activities that are similar to, but distinct from IL-12. Both IL-12 and IL-23 induce proliferation and IFN-gamma production by human T cells. While IL-12 acts on both naive and memory human T cells, the effects of IL-23 is restricted to memory T cells.

注意事项（Note）

For research use only .