## **Product Name: Recombinant Mouse LAG-3 (C-6His)**

Catalog #: PHM1056



### 概述 (Summary)

英文全称 LAG-3/CD223/Lymphocyte activation gene 3 Protein

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/μg as determined by LAL test.

蛋白构建 (Construction) Recombinant Mouse Lymphocyte Activation Gene 3 Protein is produced

by our Mammalian expression system and the target gene encoding

Ser23-Leu442 is expressed with a 6His tag at the C-terminus.

Accession # Q61790

蛋白标签 (Tag)

表达宿主 (Host) Human Cells

种属 (Species)Mouse预测分子量 (Predicted MW)46.2 KDa

蛋白形态 (Form) Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

稳定性&储存 (Stability &Storage) Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized

protein in distilled water. Please aliquot the reconstituted solution to minimize

freeze-thaw cycles.


### 电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

# **Product Name: Recombinant Mouse LAG-3 (C-6His)**

Catalog #: PHM1056





## 背景 (Background)

分子别名 (Alternative Names) 背景介绍 (References) Lymphocyte activation gene 3 protein;CD223;Lag3

Lymphocyte-activation gene 3 (LAG3), also known as CD223, is a type I transmembrane protein with four extracellular Ig-like domains, designated D1 to D4 and belongs to the immunoglobulin superfamily. The gene for LAG3 lies adjacent to the gene for CD4 on human chromosome 12p13.32 and shares approximately 20% identical to the CD4 gene. LAG3 is expressed on activated T cells, natural killer cells, B cells and plasmacytoid dendritic cells. LAG3 binds with high affinity to MHC class II molecules, and it interferes competitively with the binding of CD4 to MHC class II and thereby blocks the transduction of stimulatory signals mediated by this interaction. LAG3 negatively regulates cellular proliferation, activation, and homeostasis of T cells, and plays an important role in Treg suppressive function. LAG3 is the target of various drug development programs to develop new treatments for cancer and autoimmune disorders. The soluble form, sLAG-3, is being developed as a cancer drug.

#### 注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.