


概述 (Summary)

英文全称	Secreted Protein Acidic and Rich in Cysteine/SPARC
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Secreted Protein Acidic And Rich in Cysteine is produced by our Mammalian expression system and the target gene encoding Ala18-Ile302 is expressed with a 6His tag at the C-terminus.
Accession #	P07214
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	33.6 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse SPARC (C-6His)
Catalog #: PHM1475



背景 (Background)

分子别名 (Alternative Names)

SPARC; Sparc

背景介绍 (References)

Secreted protein acidic and rich in cysteine (SPARC, BM-40) protein belongs to the family of secreted extracellular matrix proteins with similar domain structure. Mouse SPARC protein involved an N-terminal acidic region that binds calcium, a follistatin domain containing Kazal-like sequences, and a C-terminal extracellular calcium (EC) binding domain with two EF-hand motifs. SPARC is produced by fibroblasts, capillary endothelial cells, platelets, and macrophages, especially in areas of tissue morphogenesis and remodeling. It appears to regulate cell growth through interactions with the extracellular matrix and cytokines. SPARC is expressed at high levels in tissues undergoing morphogenesis, remodeling and wound repair. The activity of SPARC is to modulate cell-cell and cell-matrix interactions, and its de-adhesive and growth inhibitory properties in non-transformed cells have led to studies to assess its role in cancer.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.