


概述 (Summary)

英文全称	Semaphorin 4A/SEMA4A
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Semaphorin 4A is produced by our Mammalian expression system and the target gene encoding Thr33-His682 is expressed with a 6His tag at the C-terminus.
Accession #	Q62178
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	72.7 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, 1mM EDTA, 5% Trehalose, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse SEMA4A (C-6His)
Catalog #: PHM1482



背景 (Background)

分子别名 (Alternative Names)

Semaphorin-4A;Semaphorin-B;Sema B;Sema4a;Semab; SemB

背景介绍 (References)

Semaphorin-4A (SEMA4A) belongs to the semaphorin family which contains a Ig-like C2-type domain, a PSI domain and a Sema domain. SEMA4A is expressed from day 10 in the embryo, and low levels are found between days 10-12. SEMA4A is a cell surface receptor for PLXNB1, PLXNB2, PLXNB3 and PLXND1 that plays an important role in cell-cell signaling. It plays a role in priming antigen-specific T-cells, promotes differentiation of Th1 T-helper cells, and thereby contributes to adaptive immunity. SEMA4A promotes phosphorylation of TIMD2, inhibits angiogenesis, and promotes axon growth cone collapse, Inhibits axonal extension by providing local signals to specify territories inaccessible for growing axons.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.