


概述 (Summary)

英文全称	Testican-3/SPOCK3
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Testican-3 is produced by our Mammalian expression system and the target gene encoding Ala23-Ile436 is expressed with a 6His tag at the C-terminus.
Accession #	Q8BKV0-1
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	47.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse Testican 3 (C-6His)
Catalog #: PHM1615



背景 (Background)

分子别名 (Alternative Names)

Testican-3;Spock3

背景介绍 (References)

Testican3, also called SPOCK3, is a secreted protein and make up a family of extracellular heparan/chondroitin sulfate proteoglycans. It contains 1 Kazal-like domain and 1 thyroglobulin type-1 domain. Testican3 contain inhibitory regions in several domains targeted to different classes of protease, and in some cases may act as protease inhibitors. In addition to their presence in testis, testicans are enriched in brain and have been shown to regulate neuronal attachment and outgrowth. It may participate in diverse steps of neurogenesis and Inhibits the processing of pro-matrix metalloproteinase 2 (MMP-2) by MT1-MMP and MT3-MMP. It also may interfere with tumor invasion.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.