


概述 (Summary)

英文全称	TFF1/Trefoil Factor 1
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Trefoil Factor 1 is produced by our Mammalian expression system and the target gene encoding Gln22-Phe87 is expressed with a 6His tag at the C-terminus.
Accession #	Q08423
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	8.3 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse TFF1 (C-6His)
Catalog #: PHM1714



背景 (Background)

分子别名 (Alternative Names)

Trefoil factor 1; Protein pS2; Tff1; Bcei; Ps2

背景介绍 (References)

Trefoil Factor 1 (TFF1) belongs to the three structurally related secreted proteins that contain trefoil domains. TFF1 is an approximately peptide that has an important effect in epithelial regeneration and wound healing. It originates from musculus and highly expressed by goblet cells of the gastric and intestinal mucosa and by conjunctival goblet cells. TFF1 is a copper-binding protein that can form disulfide-linked homodimers, associate into disulfide-linked complexes with Gastrokine 2, and form non-covalent complexes with the mucin MUC5AC. TFF1 is down-regulated during the progression from gastritis to gastric dysplasia to gastric cancer, although it is up-regulated in breast and prostate cancers. TFF1 inhibits the formation of calcium oxalate crystals, and its excretion in the urine is reduced in patients with kidney stones.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.