


概述 (Summary)

英文全称	CD40 Ligand/CD40L/TNFSF5/CD154/T-cell antigen Gp39/gp39/TNF-related activation protein/TRAP/Tumor Necrosis Factor Ligand Superfamily Member 5/CD40LG
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse CD40 Ligand is produced by our Mammalian expression system and the target gene encoding Met112-Leu260 is expressed with a 6His tag at the N-terminus.
Accession #	P27548
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	18.8 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 200mM NaCl, 0.1mM EDTA, pH 7.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse CD40L (N-6His)
Catalog #: PHM1865



背景 (Background)

分子别名 (Alternative Names)

CD40 Ligand; CD40LG;HIGM1; T-B cell-activating molecule; T-BAM; TNFSF5; tumor necrosis factor (ligand) superfamily member 5; Tumor necrosis factor ligand superfamily member 5

背景介绍 (References)

CD40 Ligand, also known as TNFSF5, CD154, is a type II transmembrane glycoprotein member of the TNF superfamily. Mature mouse CD40 Ligand consists of a 22 amino acid (aa) cytoplasmic domain, a transmembrane segment, and a 214 aa extracellular region. CD40 Ligand is expressed as a homotrimer on platelets and activated T cells and B cells. It is up/xadregulated following stimulation of basophils, eosinophils, fibroblasts, mast cells, monocytes, natural killer cells, vascular endothelial cells, and smooth muscle cells. CD40 Ligand binds and activates CD40, which is expressed on the surface of B cells, dendritic cells,macrophages, monocytes, platelets, endothelial cells, and epithelial cells. Monomeric, dimeric, and trimeric forms of soluble CD40 Ligand bind to oligomeric CD40 on cell membranes. CD40 ligation by CD40 Ligand promotes B cell activation and T cell/xaddependent humoral responses. CD40 Ligand dysregulation on T cells and antigen presenting cells contributes to the immune deficiency associated with HIV infection and AIDS. It is also implicated in the pathology of multiple cardiovascular diseases including atherosclerosis, atherothrombosis, and restenosis.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.