产品名称：Recombinant Mouse TGFBR2 (C-Fc)
产品货号：PHM2198

概述（Summary）

英文全称	TGFBR2/TGF-beta RII/TGF-beta receptor type-2/Transforming Growth Factor-β Receptor Type II (Ile24-Asp184)
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	Recombinant Mouse Transforming Growth Factor-beta Receptor Type II is produced by our Mammalian expression system and the target gene encoding Ile24-Asp184 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q62312
表达宿主（Host）	Human Cells
种属（Species）	Mouse
预测分子量（Predicted MW）	45 KDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
运输方式（Shipping）	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图（SDS-PAGE image）

产品名称：Recombinant Mouse TGFBR2 (C-Fc)
产品货号：PHM2198



背景（Background）

分子别名（Alternative Names）

TGF-beta receptor type-2; TGFR-2; TGF-beta type II receptor; Transforming growth factor-beta receptor type II; TGF-beta receptor type II; TbetaR-II; Tgfbr2

背景介绍（References）

Transforming growth factor-β (TGF-β) is an essential regulator in the processes of development, cell proliferation, and extracellular matrix deposition. TGF-β regulates cellular processes by binding to three high-affinity cell surface receptors: TGF-β receptor type I (TGF-β-RI), TGF-β receptor type II (TGF-β-RII), and TGF-ββ receptor type III (TGF-β-RIII). TGF-β RII is consists of a C-terminal protein kinase domain and an N-terminal ectodomain and belongs to transforming growth factor-beta (TGF-β) receptor subfamily. TGF-β RII has a protein kinase domain which can form a heterodimeric complex with another receptor protein and bind TGF-beta. This receptor/ligand complex phosphorylates protein will enter the nucleus and regulate the transcription of a subset of genes related to cell proliferation.

注意事项（Note）

For research use only .