产品名称：Recombinant Mouse CD7 (C-6His)
产品货号：PHM2224

概述（Summary）

英文全称	CD7/Leu-9/T-Cell Antigen CD7/GP40/T-Cell Surface Antigen Leu-9/TP41
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<0.01 EU/μg as determined by LAL test.
蛋白构建（Construction）	Recombinant Mouse T-Cell Antigen CD7 is produced by our Mammalian expression system and the target gene encoding Gln24/xadPro150 is expressed with a 6His tag at the C-terminus.
Accession #	P50283
表达宿主（Host）	Human Cells
种属（Species）	Mouse
预测分子量（Predicted MW）	15.1 KDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
运输方式（Shipping）	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图（SDS-PAGE image）

产品名称：Recombinant Mouse CD7 (C-6His)
产品货号：PHM2224



背景（Background）

分子别名（Alternative Names）

T-Cell Antigen CD7; GP40; T-Cell Leukemia Antigen; T-Cell Surface Antigen Leu-9; TP41; CD7

背景介绍（References）

T-Cell Antigen CD7 is a single-pass type I membrane protein that that belongs to the the immunoglobulin superfamily. Human CD7 is synthesized as a 240 amino acid precursor that contains a 25 amino acid signal sequence and a 215 amino acid mature chain with a Ig-like (immunoglobulin-like) domain. CD7 is normally expressed on all T-lymphocytes, NK-cells, pre-B lymphocytes and pleuripotent hematopoietic stem cells. CD7 plays an essential role in T-cell interactions, T-cell/B-cell interaction during early lymphoid development, T- and NK-cell activation and cytokine production. CD7 has been shown to interact with PIK3R1and SECTM1. However, the function of the CD7 protein in the immune system is still largely unknown.

注意事项（Note）

For research use only .