


概述 (Summary)

英文全称	CCL2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Rat C-C Motif Chemokine 2 is produced by our Mammalian expression system and the target gene encoding Gln24-Asn148 is expressed with a 6His tag at the C-terminus.
Accession #	P14844
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Rat
预测分子量 (Predicted MW)	14.9 kDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)



背景 (Background)

分子別名 (Alternative Names)

C-C motif chemokine 2; Ccl2; Monocyte chemoattractant protein 1; Monocyte chemoattractant protein 1; MCP-1; Small-inducible cytokine A2; Scya2

背景介绍 (References)

Monocyte chemoattractant protein 1 (MCP-1), also called CCL2, is a member of the beta (C-C) subfamily of chemokines that is a chemoattractant for monocytes and basophils but not eosinophils or neutrophils. CCL2 binds the cognate receptor CCR2, and together this signaling pair has been shown to have multiple pro-tumorigenic roles, from mediating tumor growth and angiogenesis to recruiting and usurping host stromal cells to support tumor progression. CCL2 is found in the circulation, where it has been suggested as a diagnostic biomarker of breast cancer and prostate cancer. CCL2 can also induce arachidonic acid release in human monocytes, which has been shown to be involved in adhesion and induction of the chemotactic response, in a pertussis-toxinsensitive manner.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.