


## 概述 (Summary)

英文全称	TGF-β2/TGF-beta 2/TGFB2/Transforming Growth Factor β-2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse/Rat Transforming Growth Factor Beta 2 is produced by our Mammalian expression system and the target gene encoding Ala303-Ser414 is expressed.
Accession #	P27090
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse/Rat
预测分子量 (Predicted MW)	12.7 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 4mM HCl.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in 4mM HCl. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in 4mM HCl. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## 电泳图 (SDS-PAGE image)



## 背景 (Background)

### 分子别名 (Alternative Names)

TGFB2; BSC-1 cell growth inhibitor; Cetermin; Glioblastoma-derived T-cell suppressor factor; G-TSF; MGC116892; Polyergin; TGF-beta2; TGF-beta-2; transforming growth factor beta-2

### 背景介绍 (References)

Transforming growth factor beta 2 (TGF- $\beta$ 2) is a member of TGF-beta superfamily that shares a characteristic cysteine knot structure. Mice with TGF- $\beta$ 2 gene deletion show defects in development of cardiac, lung, craniofacial, limb, spinal column, eye, inner ear and urogenital systems. All TGF- $\beta$  isoforms signal via the same heteromeric receptor complex, consisting of a ligand binding TGF- $\beta$  receptor type II (T $\beta$ R-II), and a TGF- $\beta$  receptor type I (T $\beta$ R-I). Signal transduction from the receptor to the nucleus is mediated via SMADs. TGF- $\beta$  expression is found in cartilage, bone, teeth, muscle, heart, blood vessels, haematopoietic cells, lung, kidney, gut, liver, eye, ear, skin, and the nervous system.

## 注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.