

Product Name: Recombinant Cynomolgus OX40L (N-8His)
Catalog #: PHV1744




概述 (Summary)

英文全称	OX40 Ligand/OX40L/TNFSF4/CD252/Tumor necrosis factor ligand superfamily member 4
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Cynomolgus OX40 Ligand is produced by our Mammalian expression system and the target gene encoding Gln51-Leu183 is expressed with a 8His tag at the N-terminus.
Accession #	F7FL80
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Cynomolgus
预测分子量 (Predicted MW)	16.5 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Cynomolgus OX40L (N-8His)
Catalog #: PHV1744



背景 (Background)

分子别名 (Alternative Names)

Tumor necrosis factor ligand 2B; TNFSF4; OX-40L

背景介绍 (References)

Tumor necrosis factor ligand superfamily member 4(TNFSF4/OX40L) is a single-pass type II membrane protein. OX40L is expressed by DCs, macrophages and B cells and signals via its cognate receptor OX40 which is mainly expressed on APCs. OX40L/OX40 interactions are important in T-cell activation and survival and for the generation of memory T cells from activated effector T cells. It is a cytokine that binds to TNFRSF4, Co-stimulates T-cell proliferation and cytokine production. OX40L/OX40 interactions are important in T-cell activation and survival and for the generation of memory T cells from activated effector T cells.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.