


Product Name: Recombinant Macaca mulatta Fetuin A (C-10His)
Catalog #: PHV1849

概述 (Summary)

英文全称	α -2-HS-Glycoprotein/AHSG/Fetuin A
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/ μ g as determined by LAL test.
蛋白构建 (Construction)	Recombinant Macaca Mulatta Alpha-2-HS-Glycoprotein is produced by our Mammalian expression system and the target gene encoding Ala19-Val367 is expressed with a 10His tag at the C-terminus.
Accession #	A0A2K6AQI2
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Macaca mulatta
预测分子量 (Predicted MW)	38.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at $\leq -70^{\circ}\text{C}$, stable for 6 months after receipt. Store at $\leq -70^{\circ}\text{C}$, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Macaca mulatta Fetuin A (C-10His)
Catalog #: PHV1849



背景 (Background)

分子别名 (Alternative Names)

Fetuin A; AHSG

背景介绍 (References)

Alpha-2-HS-glycoprotein (AHSG) is a glycoprotein that is composed of two subunits, the A and B chains, belongs to the Cystatin family of proteases inhibitors. It is highly expressed in embryonic cells and adult hepatocytes, and is expressed to a lesser extent in monocytes/macrophages. AHSG is an important circulating inhibitor of calcification in vivo, and is downregulated during the acute-phase response. It is involved in several functions, such as endocytosis, brain development and the formation of bone tissue. In addition, AHSG may influence the resolution of inflammation by modulating the phagocytosis of apoptotic cells by macrophages. ASHG blocks TGF-beta-dependent signaling in osteoblastic cells.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.