Catalog #: PHV1909



概述 (Summary)

英文全称 B7-H4/VTCN1/B7 Homolog 4/B7S1/B7x

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/µg as determined by LAL test.

蛋白构建 (Construction) Recombinant Rhesus Macaque B7-H4 is produced by our Mammalian

expression system and the target gene encoding Phe29-Ala258

expressed with a 6His tag at the C-terminus.

Accession # F7B770

蛋白标签 (Tag)

表达宿主 (Host) **Human Cells** 种属 (Species) Rhesus macaque

预测分子量 (Predicted MW) 26.4 KDa

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. 蛋白形态 (Form)

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 稳定性&储存 (Stability &Storage)

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

> Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized

> is not recommended to reconstitute to a concentration less than 100µg/ml.

protein in distilled water. Please aliquot the reconstituted solution to minimize


freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Rhesus Macaque B7-H4 (C-6His) Catalog #: PHV1909





背景 (Background)

分子别名 (Alternative Names)

B7S1; B7x; Vtcn1; B7h.5; B7-H4; B7H4T-cell costimulatory molecule B7x; B7S1VCTN1; B7XPRO1291; FLJ22418; Immune costimulatory protein B7-H4; Protein B7S1; T cell costimulatory molecule B7x; V-set domain containing T cell activation inhibitor 1; V-set domain-containing T-cell activation inhibitor 1

背景介绍 (References)

B7 Homolog 4 (B7-H4) is glycosylated member of the B7 family of immune costimulatory proteins. It is widely expressed, including in kidney, liver, lung, pancreas, placenta, prostate, spleen, testis and thymus. B7-H4 negatively regulates T-cell-mediated immune response by inhibiting T-cell activation, proliferation, cytokine production and development of cytotoxicity. When expressed on the cell surface of tumor macrophages, plays an important role, together with regulatory T-cells (Treg), in the suppression of tumor-associated antigen-specific T-cell immunity. It also involved in promoting epithelial cell transformation.

注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.