

产品名称: Recombinant Cynomolgus 4-1BB (C-6His)
产品货号: PHV1912




概述 (Summary)

| | |
|-----------------------------|---|
| 英文全称 | 4-1BB/CD137/TNFRSF9/ILA |
| 纯度 (Purity) | Greater than 95% as determined by reducing SDS-PAGE |
| 内毒素 (Endotoxin level) | <1 EU/μg as determined by LAL test. |
| 蛋白构建 (Construction) | Recombinant Cynomolgus 4-1BB ligand Receptor is produced by our Mammalian expression system and the target gene encoding Leu24-Gln186 is expressed with a 6His tag at the C-terminus. |
| Accession # | A9YYE7 |
| 表达宿主 (Host) | Human Cells |
| 种属 (Species) | Cynomolgus |
| 预测分子量 (Predicted MW) | 18.1 KDa |
| 制剂 (Form) | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4. |
| 运输方式 (Shipping) | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. |
| 稳定性&储存 (Stability &Storage) | Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months. |
| 复溶 (Reconstitution) | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

电泳图 (SDS-PAGE image)

产品名称: Recombinant Cynomolgus 4-1BB (C-6His)
产品货号: PHV1912



背景 (Background)

分子别名 (Alternative Names)

CD137; ILA; TNFRSF9; 4-1BB ligand receptor; CDw137; T-cell antigen 4-1BB homolog; T-cell antigen ILA

背景介绍 (References)

Tumor necrosis factor receptor superfamily member 9(TNFRSF9), also known as CD137 and 4-1BB, is an inducible T cell surface protein belonging to the tumor necrosis factor receptor superfamily. It is a single-pass type I membrane protein which contains 4 TNFR-Cys repeats. The human and mouse proteins share 60% amino acid sequence identity. CD137 is expressed by mesenchymal cells, including endothelial cells, chondrocytes, and cells of the central nervous system. CD137 is also broadly expressed by cells of the human immune system, is broadly expressed by cells of the human immune system, including activated CD8+ and CD4+ T cells, activated natural killer (NK) cells, follicular dendritic cells (FDCs) and monocytes. CD137 has diverse roles in the immune response, the one key function is to promote the survival of both T cells and dendritic cells by binding the cognate ligand CD137L (4-1BBL).

注意事项 (Note)

For research use only .