Product Name: Recombinant Cynomolgus VSIR (C-6His) Enkilife Catalog #: PHV2024

概述 (Summary)

英文全称 VSIR

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/μg as determined by LAL test.

蛋白构建 (Construction) Recombinant Cynomolgus V-set Immunoregulatory Receptor is produced

by our Mammalian expression system and the target gene encoding

Phe33-Ala194 is expressed with a 6His tag at the C-terminus.

Accession # A0A2K5UV12

蛋白标签 (Tag)

表达宿主 (Host)Human Cells种属 (Species)Cynomolgus预测分子量 (Predicted MW)19.2 KDa

蛋白形态 (Form) Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

稳定性&储存 (Stability &Storage) Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

is not recommended to reconstitute to a concentration less than $100\mu g/ml$. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than $100\mu g/ml$. Dissolve the lyophilized

protein in distilled water. Please aliquot the reconstituted solution to minimize


freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Cynomolgus VSIR (C-6His) Enkilife Catalog #: PHV2024





背景 (Background)

Platelet 分子别名 (Alternative Names) receptor Gi24;Stress-induced secreted protein-1;Sisp-

1;C10orf54;SISP1

背景介绍 (References) platelet receptor Gi24 is a single-pass type I membrane protein, and located at the cell surface. The protein can be cleaved by MMP14, and stimulate MMP14-mediated MMP2 activation. It is participated in the BMP signaling pathway. It also regulates the CD4-pasitive, alpha-beta T cell proliferation, and T cell cytokine production negatively. However, the protein can regulate

stem cell differentiation positively.

注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838