


Product Name: Recombinant Rhesus Macaque TNF RII (C-6His)
Catalog #: PHV2241

概述 (Summary)

英文全称	TNF RII/TNFRSF1B/CD120b/TNFR2/TNF Receptor II/Tumor Necrosis Factor Receptor II
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Rhesus Macaque Tumor Necrosis Factor Receptor II is produced by our Mammalian expression system and the target gene encoding Leu23-Asp257 is expressed with a 6His tag at the C-terminus.
Accession #	F7EAF8
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Rhesus Macaque
预测分子量 (Predicted MW)	25.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Rhesus Macaque TNF RII (C-6His)
Catalog #: PHV2241



背景 (Background)

分子别名 (Alternative Names)

Tumor necrosis factor receptor superfamily member 1b; Tnfrsf1b

背景介绍 (References)

Tumor Necrosis Factor Receptor Superfamily Member 1B (TNFRSF1B) is a member of the Tumor Necrosis Factor Receptor Superfamily. TNFRSF1B contains four TNFR-Cys repeats. TNFRSF1B can be cleaved into the following 2 chains: Tumor necrosis factor receptor superfamily member 1b and membrane form and Tumor necrosis factor-binding protein 2. TNFRSF1B is a receptor with high affinity for TNFSF2/TNF- α and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin- α . TNFRSF1B mediates most of the metabolic effects of TNF- α . TNF- α -induced apoptosis suggests that it regulates TNF- α function by antagonizing its biological activity.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.