

Product Name: Recombinant Cynomolgus CD226 (C-6His)
Catalog #: PHV2431




概述 (Summary)

英文全称	CD226
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Cynomolgus CD226 is produced by our Mammalian expression system and the target gene encoding Glu19-Asn247 is expressed with a 6His tag at the C-terminus.
Accession #	A0A2K5W1Z6
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Cynomolgus
预测分子量 (Predicted MW)	26.7 kDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Cynomolgus CD226 (C-6His)
Catalog #: PHV2431



背景 (Background)

分子别名 (Alternative Names)

CD226 molecule; DNAM-1; CD226; DNAM1

背景介绍 (References)

DNAX accessory molecule 1 (DNAM-1/CD226) is a type I transmembrane glycoprotein in the immunoglobulin superfamily. Mature human DNAM-1 contains an extracellular domain (ECD) with two Ig-like C2-set domains and a cytoplasmic region that contains motifs for binding PDZ domains and band 4.1 family proteins. DNAM-1 is expressed on multiple lymphoid and myeloid cells and interacts with CD155 and CD112. Ligation of DNAM-1 promotes the activation of NK cells, CD8+ T cells, and mast cells, dendritic cell maturation, megakaryocyte and activated platelet adhesion to vascular endothelial cells, and monocyte extravasation; it inhibits the formation of osteoclasts. Plateletendothelium, interactions mediated by DNAM-1, enable the metastasis of tumor cells to the lung.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.